

Leishmaniasis Made Ridiculously Simple



**CPT Craig G. Stark,
MD**

Walter Reed Army Medical

Leishmaniasis

- Protozoal disease
- Cutaneous leishmaniasis
 - Localized cutaneous
 - Diffuse cutaneous
 - Recidivans
 - Post-kala-azar
- Mucocutaneous leishmaniasis
- Visceral leishmaniasis
- Viscerotropic leishmaniasis



Leishmaniasis

- Parasite is found throughout the world
- Transmitted by the bite of female sandflies
- Over twenty different species
 - New World parasites
 - Old World parasites
- Uncommon in the United States
- Major epidemics worldwide



So why are we talking about this?

- Military physicians play a major role
- Incidence is rising worldwide
 - international travel
 - immigration
 - overseas military deployment
 - AIDS
- New clinical manifestations
 - Viscerotropic disease



This is demoralizing to the
troops



Background

- Ancient descriptions
- First described in English in 1756
 - “Aleppo evil”



– Other names include:

- Delhi boil, Oriental sore, Rose of Jericho, Baghdad sore, Biskra button
- Espundia, Kala-azar, black fever, dumdum fever


Background

- 1885 - Major D.D. Cunningham observed organisms from “Delhi Boil”
- 1898 - Confirmed by Russian army physician
- 1903 - British military physicians discover parasites in splenic tissue in patients with Dum-Dum Fever.
- Organism named *Leishmania donovani*



Pathophysiology

- Transmitted by sandflies
 - Promastigote stage in the sandfly
 - Amastigote stage in animal and human hosts




- Amastigotes live in macrophages
- Case reports of transmission through:
 - needle sharing, transfusions, pregnancy and sexual contact

Try to avoid this...



Pathophysiology

- Parasites incubate for weeks to months



- Varied presentation
 - immune response
 - species
 - parasite burden
- Tissue damage
- May self heal or progress to chronic disease

Frequency in the U.S.

- Uncommon
- Endemic focus in Texas
 - Southern plains woodrat
 - Coyotes and domesticated animals
 - Spread by *Lutzomyia* sandfly
- 450 military personnel
- Persian Gulf War:
 - 20 case of cutaneous disease
 - 12 cases of viscerotropic disease
- CDC reported 129 cases in 5 year period




International Frequency

- Endemic in 82 countries
- 600,000 cases per year
- 12 million infected people
- Visceral leishmaniasis
 - approximately 100,000/year
- Localized cutaneous leishmaniasis
 - 90% come from Afghanistan, Saudi Arabia, Syria, Iran and Brazil




International Frequency

- Old World disease
 - Middle East
 - Indian subcontinent
 - Southwest Asia
 - Mediterranean basin
 - East Africa
 - China
 - republics of the former soviet union
- New World disease
 - throughout the Americas
 - except Canada, Chile and Uruguay



Animal Reservoirs

- Humans are incidental hosts
- Animal reservoir
 - Wild animals
 - Domesticated dogs



Old World

- Domestic & feral dogs
- Rodents
- Foxes
- Jackals
- Wolves
- Raccoon-dogs
- Hyraxes

New World

- Sloths
- Anteaters
- Opossums
- Rodents



Human Reservoirs

- Visceral leishmaniasis
 - In India
 - human-sandfly-human
- Old World *L. tropica*
 - no animal reservoir



Morbidity/Mortality

- Cutaneous leishmaniasis
 - Localized may resolve
 - May be chronic
 - Low mortality
- Mucocutaneous leishmaniasis
 - Chronic and progressive disease
 - Variable mortality
- Visceral Leishmaniasis
 - Mortality 75-95%
 - With treatment 5% mortality



Epidemiology


- Males have increased rate of



- Male/female 2:1
- secondary to environmental exposure
 - occupation
 - leisure

Epidemiology

- Affects various age groups
 - species
 - geographic location
 - disease reservoir
 - host immune status
- Visceral leishmaniasis
 - No animal reservoir
 - All age groups
 - With animal reservoir
 - disease of childhood
- Cutaneous leishmaniasis
 - affects all ages



Localized Cutaneous Leishmaniasis

- History
 - Inoculation
 - Incubation
 - Red papule
 - No systemic symptoms
 - Localized lymphangitic spread
 - Secondary infection
 - Spontaneous healing
 - hypopigmented healed lesions
 - New world disease may progress to mucocutaneous disease



Localized Cutaneous Leishmaniasis

- Physical



- Firm papule with ulceration
- “Volcano sign”
- Fibrotic and hyperkeratotic
- Satellite lesion
- Painless
- Bacterial infection may affect presentation



Diffuse Cutaneous Leishmaniasis



- Anergic hosts
- Initial primary lesion
- Plaques and nodules
- Generally non-ulcerating
- No systemic symptoms
- Chronic
- More common in Central and South America
- May occur in East Africa

Leishmaniasis Recidivans

- Occurs after healing
- Commonly on face
- New lesions form on edges
- Infections from:
 - Reactivation
 - Infection
- Resistant to treatment



Post Kala-Azar Leishmaniasis




India

- Occurs several years after recovery
- Multiple hypopigmented dermal lesion
- Nontender
- Face and trunk
- Requires intensive treatment

Africa

- Occurs shortly after treatment
- Rash on face and extremities
- Spontaneous resolution



Mucocutaneous Leishmaniasis



- New World *L. braziliensis*
- Old World *L. aethiopica*
- Occurs after healing
- Years later mucosal involvement
- Difficult to treat
- Increased mortality
- May occur after inadequate treatment



Mucocutaneous Leishmaniasis

- Initial large lesion
- Healed scar
- Initially present with congestion
- Nare abnormalities
- “Parrot beak” or “camel nose”
- Hoarseness
- Suffocation and aspiration pneumonia



Visceral Leishmaniasis





- Fatal
- Systemic infection
- Pentad of disease:
 - fever, weight loss, hepatosplenomegaly, pancytopenia and hypergammaglobulinemia
- Night sweats, anorexia and weakness
- Characteristic skin
- “Black Fever”
- Variable incubation period
- Death from immunosuppression and secondary infection



Visceral Leishmaniasis

- Physical exam

- Cachectic patient with abdominal distension
- Liver and spleen firm and non-tender
- Epistaxis and petechia
- Lymphadenopathy
- Patchy darkening of face and trunk
- Complications



Viscerothropic Leishmaniasis

- Gulf War veterans
- Caused by *L. tropica*
- Incubation 1 month to 2 years
- Presents with nonspecific symptoms



Causes of Cutaneous Leishmaniasis

Localized cutaneous leishmaniasis:

Old World

L. donovani: China, India and Bangladesh.

L. tropica: Middle East, China, India and Mediterranean

L. major: Middle East, Africa, India, and Asia.

L. aethiopica: Ethiopia, Kenya, and Namibia.

L. infantum: Asia, Africa and Europe.

New World

L. mexicana mexicana: Central, South and North America.

L. mexicana amazonensis: Dominican Republic, Central and South America.

L. mexicana venezuelensis: Venezuela.

L. braziliensis braziliensis: Central and South America.


L. braziliensis guyanensis: Guyana, French Guyana, Surinam and Brazil.

L. braziliensis panamensis: Costa Rica, Panama, Colombia and Ecuador.

L. braziliensis peruviana: Peru and Argentina.

L. donovani chagasi: Texas, Caribbean, Central and South America.

L. shawi: South America



Causes of Cutaneous Leishmaniasis

Diffuse cutaneous leishmaniasis:

Old World

L. aethiopica: Ethiopia, Kenya and Namibia.

New World

L. mexicana mexicana: Central, South and North America.

L. mexicana amazonensis: Dominican Republic, Central/South America.

Leishmaniasis recidivans:

Old World

L. tropica: Middle East, China, India and Mediterranean.

New World

L. braziliensis braziliensis: Central and South America.

Post kala-azar leishmaniasis:


Old World

L. donovani: China, India and Bangladesh.

L. infantum: Asia, Africa and Europe.

New World

L. donovani chagasi: Central and South America



Causes of Mucocutaneous Leishmaniasis

Mucocutaneous leishmaniasis:

Old World

L. aethiopica: Ethiopia, Kenya and Namibia

New World: most often caused by

L. braziliensis braziliensis: Central and South America

L. braziliensis panamensis: Central and South America

L. braziliensis guyanensis: Guyana, Surinam and French Guiana

Also seen with

L. mexicana mexicana: Central South and North America

L. mexicana amazonensis: Brazil and Panama



Causes of Visceral Leishmaniasis

Visceral leishmaniasis:

Old World

L. donovani: China, India and Bangladesh.

L. infantum: Asia, Africa and Europe.

New World

L. donovani chagasi: Central and South America

Viscerotropic leishmaniasis:

Old World

L. tropica: Middle



Differential Diagnosis

- Cutaneous leishmaniasis:
 - Fungal- paracoccidiomycosis, chromoblastomycosis, sporotrichosis, blastomycosis
 - Bacterial- Mycobacterioses: leprosy, lupus vulgaris, tuberculosis verrucosa cutis, other mycobacterioses
 - Treponematoses- Pinta, yaws, syphilis
 - Staphylococcal/streptococcal pyodermas- impetigo, ecthyma, furunculosis
 - Parasitic- amebiasis, malaria
 - Viral- orf
 - Inflammatory- Sarcoidosis, Foreign body granuloma, pyogenic granuloma
 - Neoplastic- cutaneous T-cell lymphoma, basal cell carcinoma, squamous cell carcinoma, keratoacanthoma, cutaneous metastases
 - Granuloma faciale
 - Jessner's lymphocytic infiltrate
 - Lymphocytoma cutis
 - Discoid lupus erythematosus
 - Psoriasis
 - Keloids



Differential Diagnosis

- Mucocutaneous leishmaniasis:

- Paracoccidioidomycosis, sporotrichosis, histoplasmosis, blastomycosis, lethal midline granuloma, carcinoma, tuberculosis, gummatous syphilis, cancrum oris, yaws and rhinoscleroma

- Visceral leishmaniasis:

- Leukemia, malaria, bacterial/fungal/viral infection, tuberculosis, typhoid, and



Work Up

- History and physical
- Visualization and culture
- Montenegro Skin Test
 - similar to PPD
 - not standardized
 - made from killed promastigotes






Biopsy

- Cutaneous
 - Punch biopsy
 - Avoid necrotic center
- Mucocutaneous
 - Mucosal granuloma biopsy
 - Difficult to isolate
- Visceral
 - Bone marrow
 - Splenic aspiration
 - Contraindications
 - platelets
 - prothrombin time
 - non-palpable spleen



Biopsy

- Send for:
 - Touch preparations
 - Formalin-fixed paraffin section
- Staining:
 - Hematoxylin and eosin
 - Geimsa for touch preps and a
- Exam for:
 - Amastigote in macrophages




Culture

- Medium to
 - NNN
 - Rabbit Blood Agar
 - Schneider's Drosophila
- Positive cultures in 24 hours to weeks
- Able to culture organisms 75% of time



Additional Tests

- ELISA and direct agglutination test
- Monoclonal antibody immunofluorescence
- PCR
 - Currently being developed at WRAMC
- Rapid diagnostic antibody dipsticks
 - Quick test for visceral disease



Labs

- Cutaneous/Mucocutaneous
 - normal labs
- Visceral
 - CBC
 - Anemia
 - SPEP
 - Spike
 - LFTs
 - Mild elevations
 - Coag panel




Treatment

- Decision to treat
 - Treat visceral and mucocutaneous
 - Treat New World cutaneous disease
 - Old World cutaneous disease
 - Tends to resolve
 - Treat if:
 - Cosmetically unacceptable
 - Painful
 - Infected
 - Species dependant




Pentavalent Antimony Compounds

- Not FDA approved
- Poorly understood mechanism of action
- Two available compounds
 - Sodium Stibogluconate (Pentostam)
 - Meglumine Antimoniate (Glucantime)
- Available from the CDC and WRAMC
- Therapy
 - Expensive
 - Need prolonged treatment
 - Some side effects



Sodium Stibogluconate (Pentostam)

- Dose
 - Cutaneous: 20mg/kg IV QD for 20 days
 - Mucocutaneous: 20mg/kg IV QD for 28 days
 - Visceral: 20mg/kg IV QD for 28 days
- Side effects
 - 94% chemical pancreatitis
 - 63% elevated LFTs
 - 58% severe arthralgias/myalgias
 - 31% gastrointestinal distress
 - 27% EKG changes
- Discontinue therapy if:
 - QT interval >0.5 , Use frequent EKG monitoring
 - LFTs and amylase >5 times upper limit
 - Lipase >15 time upper limit
- Prior to therapy
 - EKG, serum electrolytes, LFTs, amylase, lipase, CBC, BUN and creatinine



Second Line Therapies

- Amphotericin B
 - Effective against resistant disease
 - Toxic side effects
 - Lipid formulation better tolerated
 - High relapse rate
- Pentamidine
 - Effective against visceral disease
 - Associated with persistent diabetes
 - High relapse rate



Second Line Therapies

- Oral medications
 - Ketoconazole, itraconazole, fluconazole, allopurinol and dapsone
- Topical
 - Paromycin effective against cutaneous disease
 - Not licensed in the U.S. but currently used by Israeli army



New Therapies

- Miltefosine

- Recent phase 2 drug study
- Showed orally administered miltefosine was 95% effective
- Phosphocholine analogue
- Treat four to six weeks
- Mild side effects :

The New England Journal of Medicine

Established in 1812 as THE NEW ENGLAND JOURNAL OF MEDICINE AND SURGERY

VOLUME 341

DECEMBER 9, 1999

NUMBER 24

ORIGINAL ARTICLES

Immediate Hormonal Therapy Compared with Observation after Radical Prostatectomy and Pelvic Lymphadenectomy in Men with Node-Positive Prostatic Carcinoma
E.M. MISSING AND OTHERS 1781

The Effect of Bisoprolol on Perioperative Mortality and Myocardial Infarction in High-Risk Patients Undergoing Vascular Surgery
D. FOLDMERMAN AND OTHERS 1789

Miltefosine, an Oral Agent, for the Treatment of Indian Visceral Leishmaniasis
T.K. JHA AND OTHERS 1795

Lack of Benefit of a Single Dose of Synthetic Human Secretin in the Treatment of Autism and Pervasive Developmental Disorders
A.D. SANDLER AND OTHERS 1801

IMAGES IN CLINICAL MEDICINE

Cystine Crystals in Fanconi's Syndrome
J.K. SINGH AND M.F. KESSOR 1807

REVIEW ARTICLES

Medical Progress: Pancreatic and Biliary Endoscopy
W.R. BRIDGER AND J. VAN DAM 1808

Mechanisms of Disease: Inflammatory Skin Diseases, T Cells, and Immune Surveillance
C. ROBERT AND T.S. KUPFER 1817

CASE RECORDS OF THE MASSACHUSETTS GENERAL HOSPITAL

A 72-Year-Old Woman with a Prosthetic Aortic Valve and Intracranial Hemorrhage
J.L. HALPERIN, S.J. FRY, AND S.L. HOOKER 1829

EDITORIALS

Early Androgen Deprivation for Prostate Cancer?
M.A. EISENBERGER AND P.C. WALSH 1837

Reducing Cardiac Risk in Noncardiac Surgery
J.H. LEI 1838

Miltefosine—The Long-Awaited Therapy for Visceral Leishmaniasis?
B.L. HERWALDT 1840

Lessons from Secretin
F.R. VORLEMAN 1842

OCASIONAL NOTES

Death of a Patient
D.M. MORENS 1845

INFORMATION FOR AUTHORS 1850

CORRESPONDENCE

Should the Declaration of Helsinki Be Revised? Aggressive Life-Lowering Therapy Compared with Angioplasty in Stable Coronary Artery Disease 1851

Late-Onset Hypertrophic Cardiomyopathy Caused by a Mutation in the Cardiac Troponin T Gene 1853

Aortic Valve Sclerosis Case 23 1999: Tuberous Pulmonary Tuberculosis Late as a Food Allergen 1846

BOOK REVIEWS 1859

NOTICES 1862

CORRECTION

Prevention of Neural Tube Defects with Folic Acid in China 1862

Owned, published, and copyrighted, 1999, by THE MASSACHUSETTS MEDICAL SOCIETY

P
E
D
I
C
A
L
S

#BXNBYC ***** 5-DIGIT 20005
#0063529/98 0000 120999 0028 27
CRIG G STORK MD
APT 1009
1305 15TH ST NW
WASHINGTON DC 20005-2951

h h

Further Care

- Prognosis
- Supportive care
- Treat bacterial infections
- Follow up in 6 weeks after the
 - Cutaneous disease
 - Visceral disease
- Retreatment with second line
 - for resistant disease
 - for reinfection



Prevention


- Vaccine trials
 - No approved vaccine
- Limit disease by
 - Reservoir eradication
 - Vector control
 - Mass treatment of patients
- Personal prevention
 - Insect repellent
 - Protective clothing
 - Permethrin-impregnated



T. Evans © 1996

Conclusions

- Increasing incidence worldwide
- Military specific disease
- Local disease may progress
- Visceral disease is fatal
- Diagnose by staining and culture
- Species ID important
- Treatment with pentavalent antimonial
- Lipid Amphotericin B second line drug
- Prevention



Questions??

